


We claim:

1. A fungicidal mixture, comprising as active components

(1) a benzamideoxime derivative of the formula I.




where the substituent and the index may have the following meanings:

R is hydrogen, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy

n is 1, 2 or 3,

and

(2) a benzophenone of the formula II,



in which

R¹ is chlorine, methyl, methoxy, acetoxy, pivaloyloxy or hydroxyl;


R² is chlorine or methyl;

R³ is hydrogen, halogen or methyl; and

R⁴ is C₁-C₆-alkyl or benzyl, where the phenyl moiety of the benzyl radical may carry a halogen or methyl substituent, and

25


(3) epoxiconazole of the formula III



10 in a synergistically effective amount.

2. A fungicidal mixture as claimed in claim 1, furthermore comprising

15 (4) pyraclostrobin of the formula IV



25 3. A fungicidal mixture as claimed in claim 1, where the radical R in the formula I is hydrogen.

4. A fungicidal mixture as claimed in any of claims 1 to 3,
30 where in the formula II

R¹ is methoxy, acetoxy or hydroxyl,

R² is methyl,

R³ is hydrogen, chlorine or bromine, and

35 R⁴ is C₁-C₄-alkyl.

5. A fungicidal mixture as claimed in claim 4, where in the formula II

40 R¹ is methoxy,

R², R⁴ are methyl and

R³ is bromine.

6. A fungicidal mixture as claimed in claim 1, where the weight ratio of the benzamideoxime derivative of the formula I to
45 the benzophenone of the formula II and the epoxiconazole of the formula III is from 20 : 1 : 1 to 1 : 20 : 20.

7. A method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with the fungicidal mixture as claimed in claim 1.
- 5 8. A method as claimed in claim 7, which comprises applying the compounds of the formulae I, II and III as set forth in claim 1 simultaneously, that is either together or separately, or in succession.
- 10 9. A method as claimed in claim 7 or 8, wherein the fungicidal mixture or the compounds of the formulae I, II and III are applied in an amount of from 0.01 to 8 kg/ha.
- 15 10. A fungicidal composition, comprising the fungicidal mixture as claimed in claim 1 and a solid or liquid carrier.

20

25

30

35

40

45


Fungicidal mixtures based on benzamideoxime derivatives,
benzophenones and an azole

5 Abstract

Fungicidal mixtures, comprising as active components

(1) a benzamideoxime derivative of the formula I

10



15

20 where the substituent and the index may have the following meanings:


R is hydrogen, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy

25

n is 1, 2 or 3,

and

30 (2) a benzophenone of the formula II,



35

in which

40 R¹ is chlorine, methyl, methoxy, acetoxy, pivaloyloxy or hydroxyl;

R² is chlorine or methyl;

R³ is hydrogen, halogen or methyl; and

45


28

R⁴ is C₁-C₆-alkyl or benzyl, where the phenyl moiety of the benzyl radical may carry a halogen or methyl substituent, and

(3) epoxiconazole of the formula III

5

10




(III)

and, if appropriate,

15

(4) pyraclostrobin of the formula IV

20



(IV)

25

in a synergistically effective amount are described.

30

35

40

45